

In Response: We agree with Montgomery et al. (1) that Chagas disease is not directly transmitted by dogs to humans. However, we emphasize that Chagas disease is in the southern United States even if risk for infection is extremely low. A study in Mexico found direct correlation of seropositivity between humans and dogs, suggesting that testing dogs may help identify prevalence of *Trypanosoma cruzi* infection among humans. They stated, "Dogs may be domestic reservoir hosts and help maintain human transmission of *T. cruzi*" (2).

For toxocariasis, indeed only embryonated eggs are infectious. In a study in the Netherlands (3), ~25% of *Toxocara* eggs found on fur were fertilized, but none were viable after 6 weeks; presence of embryonated eggs on dog fur is uncommon but can occur.

We did not mention all zoonoses that could be transmitted in a bedroom, such as toxoplasmosis or ringworm, because we could not identify

publications specifically documenting contamination in that environment. We can, however, cite examples of other infections, such as *Cheyletiella blakei* dermatitis in a woman who shared her bed with a recently acquired cat (4). We also reiterate the potential risk for human infection by the plague bacillus (*Yersinia pestis*) as a result of bed sharing, as illustrated by the case reported from Oregon in 2010 (5).

Although the risk of contracting a zoonosis in the bedroom is low, it remains possible. Bed sharing with pets should be avoided, especially for those who are immunocompromised, young, or elderly.

Bruno B. Chomel and Benjamin Sun

Author affiliations: University of California, Davis, California, USA (B.B. Chomel); and California Department of Public Health, Sacramento, California, USA (B. Sun)

DOI: 10.3201/eid1707.110590

References

1. Montgomery SP, Xiao L, Cama V. Comment on zoonoses in the bedroom [letter]. *Emerg Infect Dis* 2011;17:1340.
2. Estrada-Franco JG, Bhatia V, Diaz-Albiter H, Ochoa-Garcia L, Barbabosa A, Vazquez-Chagoyan JC, et al. Human *Trypanosoma cruzi* infection and seropositivity in dogs, Mexico. *Emerg Infect Dis*. 2006;12:624–30.
3. Overgaauw PA, van Zutphen L, Hoek D, Yaya FO, Roelfsema J, Pinelli E, et al. Zoonotic parasites in fecal samples and fur from dogs and cats in the Netherlands. *Vet Parasitol*. 2009;163:115–22. doi:10.1016/j.vetpar.2009.03.044
4. Tsianakas P, Polack B, Pinquier L, Levy Klotz B, Prost-Squarcioni C. *Cheyletiella* dermatitis: an uncommon cause of vesiculobullous eruption [in French]. *Ann Dermatol Venerol*. 2000;127:826–9.
5. Centers for Disease Control and Prevention. Notes from the field: two cases of human plague—Oregon, 2010. *MMWR Morb Mortal Wkly Rep*. 2011;60:214.

Address for correspondence: Bruno B. Chomel, Department of Population Health and Reproduction, School of Veterinary Medicine, 1114 Tupper Hall, University of California, Davis, CA 95616, USA; email: bbchomel@ucdavis.edu

Vol. 17, No. 4

References were misnumbered in the article Complete Sequence and Molecular Epidemiology of IncK Epidemic Plasmid Encoding *bla*_{CTX-M-14} (J.L. Cottell, et al.). The article has been corrected online (<http://www.cdc.gov/eid/content/17/4/645.htm>).

etymologia

Melioidosis [me"le-oi-do'sis]

From the Greek *melis*, distemper of asses, *oeidēs*, resemblance, and *osis*, a suffix indicating an abnormal condition or disease. Alfred Whitmore, a British pathologist serving in Burma, and his assistant C. S. Krishnaswami first described melioidosis in 1912. The infection became known as Whitmore's disease. In 1925, Ambrose T. Stanton and William Fletcher, the researchers who identified *Burkholderia pseudomallei* as the infection's causative agent, renamed the infection melioidosis because of its clinical resemblance to glanders.

Source: Dorland's Illustrated Medical Dictionary. 31st edition. Philadelphia: Saunders, 2007; Stanton AT, Fletcher W. Melioidosis, a disease of rodents communicable to man. *Lancet*. 1925;205:10–3. doi:10.1016/S0140-6736(01)04724-9